

**Factors Influencing the Yield
and Quality of Sorgo Sirup
Produced in West Virginia**

Summary

Performance trials of sorgo cane were conducted in West Virginia at the Ohio Valley Sub-station in 1956, 1957, and 1958. Cane and sirup yields were obtained for each year, and in 1958 sugar analyses were made on the juice from six varieties of sorgo cane harvested at two stages of maturity and held from zero to 10 days before extracting the juice. The result of these analyses indicate the following:

1. Williams and Honey yielded juices with the best sugar balances and the largest amount of finished sirup per acre, 153.7 and 151.9 gallons, respectively.
2. Growth of the cane for 10 days beyond the soft-dough stage resulted in decreased total sugar content of the juice of five of the six varieties of sorgo cane studied.
3. Holding the harvested cane for 10 days before extracting the juice resulted in complete inversion of the sucrose of the juice of five of the six varieties. Presumably the invertase content of Brawley is quite low.

THE AUTHORS

M. W. Johnson is Associate Agronomist, C. B. Sperow is Assistant Agronomist, and G. A. McLaren is Agricultural Biochemist at the West Virginia University Agricultural Experiment Station.

WEST VIRGINIA UNIVERSITY
AGRICULTURAL EXPERIMENT STATION
COLLEGE OF AGRICULTURE, FORESTRY, AND HOME ECONOMICS
A. H. VANLANDINGHAM, DIRECTOR
MORGANTOWN

Factors Influencing the Yield and Quality of Sorgo Sirup Produced in West Virginia

M. W. JOHNSON, C. B. SPEROW, and G. A. McLAREN

Introduction

THE production of sweet sorghum for sirup in West Virginia is a specialized enterprise which, although important in a few areas, is conducted only on small acreages. Most of the finished sirup produced is consumed locally, although a small amount is marketed commercially. Since information on sweet sorghum culture¹ and methods of preventing sugar crystallization had been requested by farmers of this State, a limited production performance study was begun in 1956 by the Department of Agronomy and Genetics at the Ohio Valley Sub-station at Point Pleasant.

Much of the sorgo sirup produced in West Virginia lacks consumer acceptance because of unsightly crystallization of sugars which occurs during storage. It has been reported by Walton, et al. (1938) that crystallization in sorgo sirup may result from either an excess of reducing sugars or sucrose. It is common knowledge in the field of cane sugar technology that crystallization can be prevented in high density sirups by attaining a sucrose to reducing sugar ratio of approximately one. The crystallization problem of sorgo sirups could be eliminated by producing sirups containing approximately equal percentages of sucrose and reducing sugar. This could be achieved through selection of sorgo varieties which contain the desired sucrose to reducing sugar ratio. One of the objectives of this investigation was to determine the sucrose and reducing sugar contents of six varieties of sorgo cane at different stages of maturity. A second objective involved determining the effect on the sugar composition of the juice, as the result of holding the harvested cane for 10 days before extracting the juice.

¹Detailed information on the culture of sorgo for sirup production is presented in a U.S.D.A. bulletin by Stokes, et al. (1957).

Materials and Methods

Sweet sorghum performance trials were conducted at the Ohio Valley Sub-station, Point Pleasant, in 1956, 1957, and 1958. Each year six varieties were tested. They were grown on a Wheeling fine sandy loam soil and check planted in rows spaced 42 inches apart. In 1956 and again in 1958 the plot size was 14 x 42 feet and included four rows. Thirty-five feet of the two center rows of each plot were harvested for yield. In 1957 a plot 14 x 85 feet was used with 75 feet of the two center rows harvested for yield. In all plots the varieties were planted thick and then thinned to three plants per hill. A randomized block design with four replications was used in all cases.

The trial was conducted each year on land that had been in alfalfa the previous year. Fertilizer applications consisted of 500 pounds per acre of 5-10-10 plowed down and 250 pounds applied in the row.

The weather conditions during the growing season of each of the three years varied considerably. In 1956 rainfall and temperature were average for the area; drought conditions prevailed in 1957, and in 1958 very high rainfall and lower than average temperatures occurred.

In 1956 and 1957 all of the plots were harvested when the heads of the sorgho had attained the soft-dough stage. All leaves were stripped from the stalks and the heads were removed just prior to cutting the cane. The cane was bundled, weighed, and taken to a local processing plant where the juice was extracted and processed into sirup. Because this method did not give entirely reliable estimates of finished sirup yield and quality, it was discontinued in 1958. Instead, sirup yield and quality were determined by laboratory analysis of the fresh juice. Preliminary analysis in 1957 had indicated that such a method was feasible.

The procedure followed in 1958 entailed three different harvesting treatments. In all cases the juice was extracted from a weighed sample of cane by means of a conventional cane mill. The three treatments were: (1) harvesting, extracting, and analyzing the juice of cane cut at the soft-dough stage; (2) harvesting at the soft-dough stage but extracting and analyzing the juice after a period of 10 days; and (3) harvesting, extracting, and analyzing the juice of plants cut 10 days after the soft-dough stage. A 10-pound sample of cane was used from each plot. The extracted juice was passed through a 20-mesh screen to remove any coarse foreign material and weighed. Samples of sorgho juice were frozen immediately after extraction from the cane and kept frozen until taken out for analysis.

The samples were not thawed for analysis, rather they were crushed in a hand-operated ice crusher. Fifty grams of the crushed sample was weighed on a torsion balance and immediately placed in 200 ml. of water containing 1 ml. of saturated neutral lead acetate. After addition of the lead acetate, which not only served as a clarifying agent but also inhibited plant or microbial enzyme activity, the mixture was made up to 500 ml. Solutions were permitted to stand overnight at room temperature in order to effect complete precipitation of any interfering colloids. The precipitate was removed by filtration, and excess lead was removed by means of potassium oxalate. Aliquots of the clarified, deleaded sugar solutions were analyzed for reducing sugars according to the method of Lane and Eynon, as described in the Association of Official Agricultural Chemists (1950). Sucrose was determined by difference after inversion with yeast invertase.

The sorgo juice samples were allowed to stand at room temperature for four hours before the percentage of soluble solids was determined on the supernatant. This percentage was determined by means of an Abbe refractometer equipped with a Brix scale. All percentages of soluble solids were reported as degrees Brix refractometer.

Results and Discussion

Results of these trials in terms of cane and finished sirup yields per acre for the years 1956, 1957, and 1958, are represented in Tables 1, 2, and 3, respectively. The disease factor did not present problems in any of these years so no direct statements in this regard can be made. However, certain diseases do cause heavy losses in sorgo varieties from time to time, and are an ever-present threat to their production, since all commercial varieties are susceptible to one or more of the common sorgo diseases. Detailed information on sorgo diseases is given in U.S.D.A. Farmers' Bulletin 1959 (1944).

In 1956 the only data taken were yield of cane in tons per acre and approximate yield of finished sirup in gallons per acre. The data in Table 1 indicate that yields of both cane and sirup varied considerably among the varieties tested, with sirup yields ranging from very low to very high. Environmental conditions were quite favorable for plant growth in that year. As stated previously, the method used to determine sirup yields was not entirely satisfactory and thus sirup production per acre is given as an approximation. These figures could be either somewhat higher or lower than they are as given in the tables.

In 1957 a very dry growing season was experienced at the Ohio Valley Sub-station farm, and thus the results obtained were entirely

Table 1. Sorgo Variety Trial, 1956

Variety	Yield Per Acre (Pounds)	Finished Sirup (Gallons)
Atlas -----	16,560	66.7
Williams -----	24,800	88.9
Sart -----	25,400	88.9
Tracy -----	19,400	133.3
Wiley -----	18,380	222.3
Honey -----	29,800	289.0

Table 2. Sorgo Variety Trial, 1957

Variety	Yield Per Acre (Pounds)	Finished Sirup (Gallons)
Honey -----	7,912	21.0
Sart -----	11,678	31.0
Wiley -----	10,432	41.5
Williams -----	13,273	41.5
Tracy -----	14,020	46.7
Sugar -----	19,641	62.0

different from those of 1956. Cane and sirup yields (Table 2) were greatly reduced. In several cases stands were poor, and because of the drought conditions that prevailed during July and August the plants remained in a near dormant state. Then, in September, when adequate moisture became available, growth was resumed; however, the sorgo was killed by frost before it had reached the soft-dough stage of maturity, and thus yields were drastically reduced. The variety Sugar Drip produced considerably more sirup than any of the others in this dry year. Honey, a variety that gave exceptionally high sirup yields in 1956 and 1958, produced very low cane and sirup yields in the dry year. Sirup yields were low for Sart and intermediate for Wiley, Williams, and Tracy under these same conditions.

The cane and sirup yields for the varieties tested in 1958 are given in Table 3. The finished sirup yields for this trial were calculated from cane and juice yields based on laboratory analysis and are designated as a sirup of 78 Brix (ref.). The amount of finished sirup produced per acre varied considerably among the varieties tested, but was generally quite high. Williams and Honey proved to be superior from the standpoint of both yield and quality of sirup, producing juices with suitable sucrose to reducing sugar ratios to prevent crystallization of the refined sirup (Table 2). Although the variety Sugar Drip produced a fairly high yield of finished sirup, its ratio

Table 3. Yields From Different Varieties of Sorgo Cane (1958)

Variety	Yield Per Acre (Pounds)	Juice (Per Cent)	Juice Per Acre (Pounds)	Solids Per Acre (Pounds)	Sugar Per Acre (Pounds)	Finished Sirup* (Gallons)
Brawley -----	15,326	52.2	8,153	1,003	831	91.3
Sart -----	20,140	50.6	10,191	1,070	895	98.4
Tracy -----	15,664	52.9	8,286	1,243	960	105.5
Sugar -----	16,908	58.3	9,857	1,311	1,112	122.2
Honey -----	16,420	61.8	10,148	1,563	1,382	151.9
Williams -----	20,945	56.4	11,813	1,583	1,399	153.7

*The term finished sirup is used to designate a sirup of 78 Brix (ref.).

of sucrose to reducing sugar was completely out of balance, as was that of the other lower yielding varieties included in the trial. The sucrose content of Sart, Brawley, and Sugar Drip was too high to be certain that crystallization would not occur. Although partial inversion of sucrose from the juice of these sorgo varieties would yield a sirup not too prone to crystallization, such a step must be well-controlled and would increase the cost of production. The high reducing sugar content of juice extracted from the Tracy variety indicates that sirups produced from such juice would show crystallization of reducing sugars.

Cutting the cane 10 days after it reached the soft-dough stage did not markedly affect the ratio of sucrose to reducing sugar, but it did influence the percentage of total sugars in the juice (Table 4). There was a decrease in total sugars in the juice of all varieties except Tracy, which showed a slight increase in total sugars.

It may be noted at the bottom of Table 4 that holding cane for 10 days after harvesting resulted in complete inversion of the sucrose in the juice from all varieties except Brawley. Apparently all of the other varieties contained sufficient invertase to effect the inversion of sucrose to invert sugar (reducing sugars). These data certainly indicate that cane should be extracted immediately after it is cut, rather than held for several weeks before extraction. Although it is a common practice of many sorgo producers to hold cane after it is cut in hope that the resulting inversion will prevent crystallization, actually it appears to be responsible for most of the crystallization problems. There is no way of controlling inversion of sucrose by the invertase in the cane, hence a variety containing a high concentration of this enzyme will completely invert all of the sugar just as happened in five of the varieties used in this study.

Literature Cited

- A.O.A.C. 1950. *Methods of Analysis of the Association of Official Agricultural Chemists*. Washington, D. C.
- Leukel, R. W., John H. Martin, and C. L. Leefebore. 1944. *Sorghum Diseases and Their Control*. Farmers' Bulletin No. 1959., U.S.D.A.
- Stokes, I. E., O. H. Coleman, and Jack L. Dean. 1957. *Culture of Sorgo for Sirup Production*. Farmers' Bulletin No. 2100., U.S.D.A.
- Walton, C. F., E. K. Ventre, and S. Bell. 1938. *Farm Production of Sorgo Sirup*. Farmers' Bulletin No. 1791., U.S.D.A.

Table 4. The Effect of Stage of Maturity of Cane on the Sugar Composition of Sorgo Juice (1958)

Variety	Stage of Maturity of Cane When Cut	Time of Extraction of the Cane	Number of Plots	Percentage of Sugars on a Dry Matter Basis			Soluble Solids Brix (ref.) Deg.
				Reducing Sugars	Sucrose	Total Sugars	
Brawley	Soft-dough	Immediately after cutting	4	25.5	57.4	82.9	12.3
Tracy	"	"	2	57.9	19.3	77.2	15.0
Sart	"	"	2	25.7	57.9	83.6	10.5
Williams	"	"	2	47.3	41.1	88.4	13.4
Honey	"	"	2	48.2	37.9	86.1	15.4
Sugar	"	"	2	26.9	57.9	84.8	13.3
Brawley	Soft-dough + 10 days	"	4	13.2	68.8	81.9	12.1
Tracy	"	"	2	51.0	30.7	81.7	12.6
Sart	"	"	2	45.7	30.5	76.2	9.0
Williams	"	"	2	37.5	35.7	73.2	12.2
Honey	"	"	2	36.9	35.8	72.7	11.8
Sugar	"	"	2	26.3	51.7	78.0	10.0
Brawley	Soft-dough	10-days after cutting	4	33.9	44.2	78.1	17.3
Tracy	"	"	2	Sucrose completely inverted to reducing sugars			
Sart	"	"	2	"	"	"	"
Williams	"	"	2	"	"	"	"
Honey	"	"	2	"	"	"	"
Sugar	"	"	2	"	"	"	"